



KJM 9250

$^1\text{H}$  NMR spectra on the AVI-600 and AVII-600

Version 1.0

Topspin 3.5 Windows 7  
Topspin 1.3 Windows XP



© Professor Emeritus Alistair Lawrence Wilkins,  
University of Waikato, New Zealand.  
January 2018

# **<sup>1</sup>H NMR spectra on the AVI-600 and AVII-600**

## **1.0 Introduction**

aw coded <sup>1</sup>H NMR parameter files generally use a 90° pulse for maximum <sup>1</sup>H signal.

Best <sup>1</sup>H resolution is obtained using **FT** and **PK** (or **APK**) processing. **FT** processing does not apply a line broadening factor. **EF** or **EFP** processing applies a line broadening factor (**LB**).

Resolution enhancement uses negative **LB** values. Try **LB** = -1.0 to -2.5 Hz with **GB** = 0.33, and **GFP** processing. Remember to reset **LB** and **GB** to their normal values (0.1 and 0 respectively) after **GFP** processing.

## **1.1 Presaturation Experiments**

Continuous wave or excitation sculptured (ES) can be used to presaturate <sup>1</sup>H NMR signals. The simplest of these techniques is continuous wave presaturation.

**CW** presaturation power levels (db settings) can be increased or decreased by subtracting or adding 3-12 db respectively. 6 db = a factor of 2.

The **ES** shaped pulse's excitation window can be decreased by doubling its shaped pulse time from 2000 usec to 4000 usec and halving its power by adding 6 db to that read in using the **getprosol** command.

## **2.0 <sup>1</sup>H NMR experiments**

**2.1 <sup>1</sup>H NMR with a 30, 45 or 90 degree pulse**

**2.2 <sup>1</sup>H NMR with CW presaturation**

**2.3 <sup>1</sup>H NMR with dual CW presaturation**

**2.4 <sup>1</sup>H NMR with ES peak suppression**

**2.5 <sup>1</sup>H NMR with combined ES + CW presaturation on F1**

**2.6 <sup>1</sup>H NMR with combined ES + CW presaturation on F1  
and CW presaturation on F2**

**2.7 <sup>1</sup>H NMR with three peak ES + dual CW presaturation**

## 2.1 $^1\text{H}$ NMR spectra with a 30, 45 or 90 degree pulse

Parameter sets: **awproton30**, **awproton45**, **awproton90** (+ **getprosol**)

Pulse programmes: **zg30**, **awzg45** or **zg** respectively

**TD** = 64 K, **SI** = 64 K

**SW** = 16 ppm, **O1P** = 7.0 ppm.


**D1** = 1.5 sec or other time of your choice.

**NS** = any number, **DS** = 2, 4, 8.

Type **ased** (enter) and review parameters used in the job.

Set **receiver gain** using **RGA** (*important!*).

Process with **FT** (no line broadening) or **EFP** (applies **LB**).



600 MHz  $^1\text{H}$  NMR spectrum of quinine in  $\text{D}_6$ -DMSO.

## 2.2 $^1\text{H}$ NMR spectrum with CW presaturation

Parameter set: **awprotonpr** (+ **getprosol**)

Pulse programme: **zgpr**

**TD** = 32 K or 64 K , **SI** = 32 K or 64 K.

**SW** = 18 ppm.

**O1** = frequency in Hz of the F1 signal to be presaturated.

= spectral window midpoint. Check **SW** is wide enough.

**PL9** = **F1** presaturation power applied during **D1**.

**D1** = 2 sec or other time of your choice.


Type **ased** (enter) and review parameters used in the job.

Add (or subtract) 3-12 db to **PL9** to decrease (or increase) the presaturation power.

6 db = a factor of 2. A larger attenuation setting decreases the power level.

Set **receiver gain** using **RGA** (*important!*).

Process with **EFP** (applies **LB**).



**Lower:** 600 MHz  $^1\text{H}$  NMR spectrum of quinine in  $\text{D}_6$ -DMSO.

**Upper:**  $^1\text{H}$  NMR spectrum with CW presaturation of the HOD line at 3.37 ppm.

### 2.3 $^1\text{H}$ NMR spectrum with dual CW presaturation

Parameter set: **awprotonprf1prf2 (+ getprosol)**

Pulse programme: **awprotonprf1prf2**

**TD** = 64 K, **SI** = 64 K.

**SW** = 18 ppm.

**O1** = frequency in Hz of the F1 signal to be presaturated.  
= spectral window midpoint. Check **SW** is wide enough.

**O2** = frequency in Hz of the F2 signal to be presaturated.

**PL9** = F1 presaturation power applied during D1.

**PL21** = F2 presaturation power applied during D1.


**D1** = 2 sec or other time of your choice.

Type **ased** (enter) and review parameters used in the job.

Add (or subtract) 3-12 db to **PL9** and/or **PL21** to decrease (or increase) the presaturation power. 6 db = a factor of 2. A larger attenuation setting decreases the power level.

Set **receiver gain** using **RGA** (*important!*).

Process with **EFP** (applies **LB**).



**Lower:** 600 MHz  $^1\text{H}$  NMR spectrum of quinine in  $\text{D}_6$ -DMSO.

**Upper:**  $^1\text{H}$  NMR spectrum with CW presaturation of the HOD (3.37 ppm) and DMSO (2.5 ppm) lines.

## 2.4 $^1\text{H}$ NMR spectrum with ES peak suppression

Parameter sets: **awprotones (+ getprosol)**

Pulse programmes: **zgesgp**

**TD** = 64 K, **SI** = 64 K.

**SW** = 18 ppm.

**O1** = frequency in Hz of the F1 signal to be ES suppressed.

= spectral window midpoint Check. **SW** is wide enough.


**D1** = 2 sec or other time of your choice.

Type **ased** (enter) and review parameters used in the job. Verify gradients are OK.

Check **P12** = 2000 usec, **SPNAM1** = **squa100.1000**

Set **receiver gain** using **RGA** (*important!*).

Process with **EFP** (applies **LB**).



**Lower:** 600 MHz  $^1\text{H}$  NMR spectrum of quinine in  $\text{D}_6\text{-DMSO}$ .

**Upper:**  $^1\text{H}$  NMR spectrum with ES suppression of the HOD line at 3.37 ppm.

## 2.5 $^1\text{H}$ NMR with combined ES and CW presaturation on F1

Parameter set: **awprotonespr** (+ **getprosol**)

Pulse programmes: **awprotonespr**

**TD** = 64 K, **SI** = 64 K.

**SW** = 18 ppm.

**O1** = frequency in Hz of the F1 signal to be ES suppressed.  
= spectral window midpoint. Check **SW** is wide enough.


**PL9** = F1 presaturation power applied during D1.

**D1** = 2 sec or other time of your choice.

Type **ased** (enter) and review parameters used in the job. Verify gradients are OK.  
Check **P12** = 2000 usec, **SPNAM1** = **squa100.1000**.

Set **receiver gain** using **RGA** (*important!*).

Process with **EFP** (applies **LB**).



**Lower:** 600 MHz  $^1\text{H}$  NMR spectrum of quinine in  $\text{D}_6$ -DMSO.

**Upper:**  $^1\text{H}$  NMR spectrum with combined ES and CW presaturation of the HOD line at 3.37 ppm.

## 2.6 $^1\text{H}$ NMR spectrum with combined ES+CW presaturation on F1 and CW presaturation on F2

Parameter set: **awprotonesprf1prf2 (+ getprosol)**

Pulse programmes: **awprotonesprf1prf2**

**TD** = 64 K, **SI** = 64 K.

**SW** = 18 ppm.

**O1** = frequency in Hz of the F1 signal to be combined ES + CW suppressed  
= spectral window mid-point. Check SW is wide enough..

**O2** = frequency in Hz of the F2 signal to be CW presaturated.

**PL9** = F1 presaturation power applied during **D1**.

**PL21** = F2 presaturation power applied during **D1**.


**D1** = 2 sec or other time of your choice.

Type **ased** (enter) and review parameters used in the job. Verify gradients are OK.

Check **P40** = 2000 usec, **SPNAM10** = **squa100.1000**.

Set **receiver gain** using **RGA** (*important!*).

Process with **EFP** (applies **LB**).



**Lower:** 600 MHz  $^1\text{H}$  NMR spectrum of quinine in  $\text{D}_6$ -DMSO.

**Upper:**  $^1\text{H}$  NMR with combined ES + CW presaturation of the HOD line (3.37 ppm) on F1 and the DMSO line (2.5 ppm) on F2.

## 2.7 $^1\text{H}$ NMR spectrum with three peak ES + dual CW presaturation

Parameter set: **awprotonesprf1prf2** (+ **getprosol**)

Pulse programmes: **awprotonesprf1prf2**

**TD** = 64 K, **SI** = 64 K.

**SW** = 20 ppm.

**O1** = frequency in Hz of the F1 signal to be CW suppressed.

= spectral window midpoint. Check SW is wide enough.

**O1\*** = frequency in Hz of the F1 signal to be ES suppressed

**SPOFFS10** = (**O1\***-**O1**) Hz (may be a positive or negative value).

**O2** = frequency in Hz of the F2 signal to be CW presaturated.

**PL9** = F1 presaturation power applied during **D1**.

**PL21** = F2 presaturation power applied during **D1**.


**D1** = 2 sec or other time of your choice.

Type **ased** (enter) and review parameters used in the job. Verify gradients are OK.

Check **P40** = 2000 usec, **SPNAM10** = **squa100.1000**.

Set **receiver gain** using **RGA** (*important!*).

Process with **EFP** (applies **LB**).



**Lower:** 600 MHz  $^1\text{H}$  NMR spectrum of quinine in  $\text{D}_6$ -DMSO.

**Upper:**  $^1\text{H}$  NMR with CW presaturation on F1 of quinine's  $\text{OCH}_3$  signal (3.89 ppm), offset ES suppression of the HOD line (3.37 ppm) and CW presaturation on F2 of the DMSO signal (2.5 ppm).