

The fractional Ornstein-Uhlenbeck process (fOU)

The fOU process with forcing term (ffOU)

The time-changed fractional Ornstein-Uhlenbeck process (TCfOU)

The limit distribution of the TCfOU process

Convergence results for the TCfOU process

On a fractional Ornstein-Uhlenbeck process and its time-changed version

*Some results obtained with
Yuliya Mishura and Giacomo Ascione*

Enrica Pirozzi

Dipartimento di Matematica e Applicazioni,
Università di Napoli Federico II, Napoli, Italy

The fractional Ornstein-Uhlenbeck process (fOU)

The fOU process with forcing term (ffOU)

The time-changed fractional Ornstein-Uhlenbeck process (TCfOU)

The limit distribution of the TCfOU process

Convergence results for the TCfOU process

Outline

1 The fractional Ornstein-Uhlenbeck process (fOU)

The fractional Ornstein-Uhlenbeck process (fOU)

The fOU process with forcing term (ffOU)

The time-changed fractional Ornstein-Uhlenbeck process (TCfOU)

The limit distribution of the TCfOU process

Convergence results for the TCfOU process

Outline

- 1 The fractional Ornstein-Uhlenbeck process (fOU)
- 2 The fOU process with forcing term (ffOU)
 - The mean value function
 - The covariance function
 - Covariance as the function of time and Hurst index

The fractional Ornstein-Uhlenbeck process (fOU)

The fOU process with forcing term (ffOU)

The time-changed fractional Ornstein-Uhlenbeck process (TCfOU)

The limit distribution of the TCfOU process

Convergence results for the TCfOU process

Outline

- 1 The fractional Ornstein-Uhlenbeck process (fOU)
- 2 The fOU process with forcing term (ffOU)
 - The mean value function
 - The covariance function
 - Covariance as the function of time and Hurst index
- 3 The time-changed fractional Ornstein-Uhlenbeck process (TCfOU)
 - The time-change process

The fractional Ornstein-Uhlenbeck process (fOU)

The fOU process with forcing term (ffOU)

The time-changed fractional Ornstein-Uhlenbeck process (TCfOU)

The limit distribution of the TCfOU process

Convergence results for the TCfOU process

Outline

- 1 The fractional Ornstein-Uhlenbeck process (fOU)
- 2 The fOU process with forcing term (ffOU)
 - The mean value function
 - The covariance function
 - Covariance as the function of time and Hurst index
- 3 The time-changed fractional Ornstein-Uhlenbeck process (TCfOU)
 - The time-change process
- 4 The limit distribution of the TCfOU process

Outline

- 1 The fractional Ornstein-Uhlenbeck process (fOU)
- 2 The fOU process with forcing term (ffOU)
 - The mean value function
 - The covariance function
 - Covariance as the function of time and Hurst index
- 3 The time-changed fractional Ornstein-Uhlenbeck process (TCfOU)
 - The time-change process
- 4 The limit distribution of the TCfOU process
- 5 Convergence results for the TCfOU process
 - Weak one-dimensional convergence of the TCfOU process
 - A functional limit theorem for the TCfOU process

The fractional Ornstein-Uhlenbeck process

It is the solution process of the following linear stochastic differential equation:

$$dV_t = -\frac{1}{\theta} V_t + \sigma dB_t^H, V_0 = \xi \quad (1)$$

in a complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$,
 where $B^H = \{B_t^H, t \geq 0\}$ is a fractional Brownian motion (fBm)
 with Hurst parameter $H \in (0, \frac{1}{2}) \cup (\frac{1}{2}, 1)$,
 and $\mathcal{F}_t^H, t \geq 0$ be its natural filtration.

Specifically, the fBm is the almost surely path-continuous centered Gaussian process $B^H = \{B_t^H, t \geq 0\}$ with covariance function given by

$$\mathbb{E}[B_t^H B_s^H] = \frac{1}{2}(|t|^{2H} + |s|^{2H} - |t - s|^{2H}).$$

At first, motivated by the wish to construct a new fractional neuronal model, we consider the following linear stochastic differential equation driven by our fBm:

$$dV_t = \left[-\frac{1}{\theta}(V_t - \tilde{V}) + I_t \right] dt + \sigma dB_t^H, \quad V_0 = \xi \quad (2)$$

where ξ is a square-integrable random variable defined on $(\Omega, \mathcal{F}, \mathbb{P})$, $I_t = I$ is a forcing term, $\theta > 0$ and $\tilde{V}$ are constant.

The above equation is equivalent to the integral equation

$$V_t = \xi - \frac{1}{\theta} \int_0^t (V_s - \tilde{V}) ds + \int_0^t I_s ds + \sigma B_t^H. \quad (3)$$

We call the solution $V = V_t$ of equation (2) ffOU process with forcing term $I = I_t$ and $\tilde{V}$ its resting term, that is to say a globally asymptotically stable equilibrium for the expected value as $I_t \equiv 0$.

Referring to Kaarakka T, Salminen P (2011), and in particular, following the lines of proposition A.1 of Cheridito et al. (2003) , we can explicitly determine the solution (denoting $\mathcal{F}_t^{H,\xi}$ the sigma-field generated by ξ and $B_s^H, s \leq t$).

Proposition

If I is a stochastic process such that its sample paths are (almost surely) integrable in any interval $[0, T]$ for $T > 0$, then equation (3) admits a unique solution whose paths are almost surely continuous. This solution can be expressed as

$$V_t = \tilde{V} + e^{-\frac{t}{\theta}} \left(-\tilde{V} + \xi + \int_0^t I_s e^{\frac{s}{\theta}} ds + \sigma \int_0^t e^{\frac{s}{\theta}} dB_s^H \right), \quad (4)$$

where the integral is point-wise interpreted as a Riemann-Stieltjes integral ([Biagini et al, 2008]). In the case $H < 1/2$ it is defined via integration by parts, namely,

$$V_t = \tilde{V} + e^{-\frac{t}{\theta}} \left(-\tilde{V} + \xi + \int_0^t I_s e^{\frac{s}{\theta}} ds + \sigma \left(e^{\frac{t}{\theta}} B_t^H - \theta^{-1} \int_0^t e^{\frac{s}{\theta}} B_s^H ds \right) \right). \quad (5)$$

Moreover, V_t is $\mathcal{F}_t^{H,\xi}$ -adapted if I_t is a $\mathcal{F}_t^{H,\xi}$ -adapted process on $(\Omega, \mathcal{F}, \mathbb{P})$.

The mean value function

Recalling that

$$\mathbb{E} \left(\int_0^t e^{\frac{s}{\theta}} dB_s^H \right) = 0$$

for any $t > 0$, we immediately get the following result.

Proposition

Let I be a stochastic process defined on $(\Omega, \mathcal{F}, \mathbb{P})$ such that for any $t > 0$, $I_t \in L^1(\Omega, \mathcal{F}, \mathbb{P})$, and $\mathbb{E}[|I_t|] \in L^1([0, T], \lambda)$, where λ is the Lebesgue measure. Also, let $V = V_t$ be the solution of equation (2). Then

$$\mathbb{E}[V_t] = (1 - e^{-\frac{t}{\theta}}) \tilde{V} + e^{-\frac{t}{\theta}} \mathbb{E}[\xi] + e^{-\frac{t}{\theta}} \int_0^t e^{\frac{s}{\theta}} \mathbb{E}[I_s] ds. \quad (6)$$

The covariance function

In order to exploit the results of [Cheridito et al., 2003], we introduce the **stationary** fractional Ornstein-Uhlenbeck process U^H with parameter $\frac{1}{\theta}$ (sfOU for short), as the only stationary Gaussian solution of equation (7)

$$dU_t^H = -\frac{1}{\theta} U_t^H dt + \sigma dB_t^H. \quad (7)$$

with the initial value

$$U_0^{H,\eta} = \eta$$

where η is the following square integrable random variable

$$\eta = \sigma \int_{-\infty}^0 e^{\frac{s}{\theta}} dB_s^H$$

We finally refer to the process

$$U_t^{H,\eta} = e^{-\frac{t}{\theta}} \left(\eta + \sigma \int_0^t e^{\frac{s}{\theta}} dB_s^H \right) = \sigma \int_{-\infty}^t e^{-\frac{t-s}{\theta}} dB_s^H = U_t^H, \quad t \geq 0$$

which is a stationary long-range dependent process for $H > \frac{1}{2}$,
while is short-range dependent for $H < \frac{1}{2}$.

A closed form of its covariance function for $H \in (0, 1)$ was obtained in [Pipiras and Taqqu, 2000], (see also [Cheridito et al., 2003]), Remark 2.4). It has a form

$$\rho(s) := \text{Cov}(U_t^H, U_{t+s}^H) = \sigma^2 \theta^2 C_H \int_{-\infty}^{+\infty} \frac{|x|^{1-2H}}{1 + \theta^2 x^2} e^{isx} dx, \quad (8)$$

where

$$C_H = \frac{\Gamma(2H + 1) \sin(\pi H)}{2\pi}. \quad (9)$$

It is also shown in [Cheridito et al., 2003] that, for $s \rightarrow \infty$, the covariance function admits the following asymptotic expression for any $H \in (0, 1) \setminus \{1/2\}$, $N = 1, 2, \dots$ and for fixed t :

$$\text{Cov}(U_t^H, U_{t+s}^H) = \frac{1}{2} \sigma^2 \sum_{n=1}^N \theta^{2n} \left(\prod_{k=0}^{2n-1} (2H - k) \right) s^{2H-2n} + O(s^{2H-2N-2}).$$

In particular, for $N = 1$, we obtain

$$\text{Cov}(U_t^H, U_{t+s}^H) = \sigma^2 \theta^2 H(2H - 1) s^{2H-2} + O(s^{2H-4}). \quad (10)$$

For $\frac{1}{2} < H < 1$, we have $-1 < 2H - 2 < 0$, then

$$\text{Cov}(U_t^H, U_{t+s}^H) \sim K s^{2H-2}$$

as $s \rightarrow +\infty$, for $K = \sigma^2 \theta^2 H(2H - 1)$, saying, as usual, that $f \sim g$ if $\lim_{s \rightarrow +\infty} \frac{f(s)}{g(s)} = 1$.

Hence the long-range dependence of the process U^H for $H > \frac{1}{2}$ follows.

Conversely, for $0 < H < \frac{1}{2}$ we have $-2 < 2H - 2 < -1$ whence the short-range dependence follows.

Using these results and Theorem 2.3 in [Cheridito et al., 2003] we can get the covariance function in the following symmetric form.

Lemma

Let $U^{H,x} = U_t^{H,x}$, $t \geq 0$ be a fOU process solving equation (7) with initial value $x \in \mathbb{R}$. Then its covariance does not depend on $x \in \mathbb{R}$ and has a form

$$\begin{aligned} R_H(t, s) &:= \text{Cov}(U_t^{H,x}, U_s^{H,x}) \\ &= \sigma^2 \theta^2 C_H \int_{\mathbb{R}} \left(e^{isy} - e^{-\frac{s}{\theta}} \right) \left(e^{ity} - e^{-\frac{t}{\theta}} \right) \frac{|y|^{1-2H}}{1 + \theta^2 y^2} dy. \end{aligned} \quad (11)$$

The proof is essentially based on

$$U_t^{H,x} - \mathbb{E}[U_t^{H,x}] = \sigma \int_0^t e^{-\frac{t-v}{\theta}} dB_v^H,$$

then, we have for $s \geq t$

$$\begin{aligned} R_H(t, s) &= \sigma^2 \mathbb{E} \left[\left(\int_0^t e^{-\frac{t-v}{\theta}} dB_v^H \right) \left(\int_0^s e^{-\frac{s-u}{\theta}} dB_u^H \right) \right] \\ &= \sigma^2 \left(\mathbb{E} \left[\left(\int_{-\infty}^t e^{-\frac{t-v}{\theta}} dB_v^H - \int_{-\infty}^0 e^{-\frac{t-v}{\theta}} dB_v^H \right) \right. \right. \\ &\quad \times \left. \left(\int_{-\infty}^s e^{-\frac{s-u}{\theta}} dB_u^H - \int_{-\infty}^0 e^{-\frac{s-u}{\theta}} dB_u^H \right) \right] \right) \\ &= \rho(s-t) - e^{-\frac{t}{\theta}} \rho(s) - e^{-\frac{s}{\theta}} \rho(t) + e^{-\frac{t+s}{\theta}} \rho(0). \end{aligned} \tag{12}$$

Covariance function of ffOU

Now we proceed with the covariance function for ffOU process, i.e., fOU process with non-zero forcing term I . Denote

$$c(u, v) = \text{Cov}(I_u, I_v)$$

From now on, we assume that the initial value $\xi = x \in \mathbb{R}$.

Lemma

With the notation above specified, let ξ be a degenerate random variable. Suppose I is a stochastic process defined on $(\Omega, \mathcal{F}, \mathbb{P})$ such that:

- (i) For any $t > 0$, $I_t \in L^2(\Omega, \mathcal{F}, \mathbb{P})$.
- (ii) For any $T > 0$, $t \mapsto \mathbb{E}[|I_t|]$ is integrable in $[0, T]$.
- (iii) For any $(t, s) \in \mathbb{R}^2$ such that $t, s > 0$, $(u, v) \mapsto c(u, v)$ is integrable in $[0, t] \times [0, s]$.
- (iv) For any $t, s > 0$ random variables I_t and B_s^H are uncorrelated.

Then

$$\text{Cov}(V_t, V_s) = e^{-\frac{t+s}{\theta}} \iint_{[0,t] \times [0,s]} e^{\frac{u+v}{\theta}} c(u, v) du dv + R_H(t, s). \quad (13)$$

Covariance as the function of time and Hurst index.

Asymptotic behavior of covariance

Non-random forcing term and consider $R_H(t, t + s)$.

To start, note that in [Cheridito et al., 2003] the authors proved the following asymptotic expansion for fixed $t \geq 0$, $N = 1, 2, \dots$ and $s \rightarrow +\infty$:

$$\begin{aligned}
 R_H(t, t + s) = & \\
 & \frac{1}{2} \sigma^2 \sum_{n=1}^N \theta^{2n} \left(\prod_{k=0}^{2n-1} (2H - k) \right) \left[s^{2H-2n} - e^{-\frac{t}{\theta}} (t + s)^{2H-2n} \right] \\
 & + O(s^{2H-2N-2}),
 \end{aligned}$$

which, for $N = 1$, becomes

$$R_H(t, t+s) = \sigma^2 \theta^2 H(2H-1)(s^{2H-2} - e^{-\frac{t}{\theta}}(t+s)^{2H-2}) + O(s^{2H-4})$$

so that also

$$R_H(t, t+s) \sim K_t s^{2H-2}$$

with a constant

$$K_t = \sigma^2 \theta^2 H(2H-1)(1 - e^{-t})$$

For this reason we can conclude that $U_t^{H,x}$, as well as U_t^H , demonstrates a time non-homogeneous **long-range dependence** for $H > \frac{1}{2}$ and a time non-homogeneous **short-range dependence** for $H < \frac{1}{2}$.

The fractional Ornstein-Uhlenbeck process (fOU)

The fOU process with forcing term (ffOU)

The time-changed fractional Ornstein-Uhlenbeck process (TCfOU)

The limit distribution of the TCfOU process

Convergence results for the TCfOU process

The mean value function

The covariance function

Covariance as the function of time and Hurst index




Figure: The covariance function $R_H(t, t+s)$ for $\theta = 30$, $\sigma = 1$, $t = 10$ for a non-random I : on the left as a function of s for different values of H ; on the right as a function of H for different values of s .

Figure 1 on the left demonstrates that the tails of $s \mapsto R_H(t, t+s)$ as $s \rightarrow \infty$ depend on the Hurst parameter and they are slower in convergence to 0 as H grows.

Random Forcing Term

Now, let I be stochastic. Denote $C_V(t, s) = \text{Cov}(V_t, V_s)$.

In general, it is not obvious that $C_V(t, s) \rightarrow 0$ as $s \rightarrow +\infty$.

The following lemma establishes some hypotheses under which $C_V(t, s) \rightarrow 0$ as $s \rightarrow +\infty$.

Lemma

Suppose the forcing term I verifies the hypotheses of previous Lemma. Then $\lim_{s \rightarrow +\infty} C_V(t, s) = 0$ under one of the following additional hypothesis:

$\mathcal{L}1$ The function $v \mapsto \int_0^t e^{\frac{u+v}{\theta}} c_I(u, v) du$ is in $L^1(\mathbb{R}^+)$;

$\mathcal{L}2$ The following properties are verified:

(i)

$$(ii) \quad \lim_{s \rightarrow +\infty} \int_0^s \int_0^t e^{\frac{u+v}{\theta}} c_I(u, v) dudv = \infty;$$

$$\lim_{s \rightarrow +\infty} \int_0^t e^{\frac{u}{\theta}} c_I(u, s) du = 0.$$

Moreover, property (ii) of hypothesis $\mathcal{L}2$ is assured if the following properties hold:

(iii) There exists $k \in L^1([0, t])$ such that $|c_I(u, s)| \leq k(u)$ for almost all $u \in [0, t]$ and $s \geq 0$.

(iv) $\lim_{s \rightarrow +\infty} c_I(u, s) = 0$.

Theorem

Suppose the forcing term I verifies the hypotheses of previous Lemma. Then, $C_V(t, s) \sim Ks^{2H-2}$ for some constant K under one of the following additional hypothesis: hypothesis $\mathcal{L}1$ from Lemma 3 or

$\mathcal{L}2'$ The following properties are verified: property (i) from hypothesis $\mathcal{L}2$ from Lemma 3 and
(ii)'

$$\lim_{s \rightarrow +\infty} s^{2-2H} \int_0^t e^{\frac{u}{\theta}} c_I(u, s) du = 0.$$

Moreover, property (ii)' of hypothesis $\mathcal{L}2'$ is assured by the following properties:

- (iii)' there exists $k \in L^1([0, t])$ such that $s^{2-2H} |c_I(u, s)| \leq k(u)$ for almost all $u \in [0, t]$ and $s \geq 0$;
- (iv)' $\lim_{s \rightarrow +\infty} s^{2-2H} c_I(u, s) = 0$.

By this theorem, we know that under hypothesis $\mathcal{L}1$ or $\mathcal{L}2'$,

$$C(t, s) \sim K_t s^{2H-2}$$

Hence, in such case V_t exhibits a time non-homogeneous long-range dependence for $H > \frac{1}{2}$ and a time non-homogeneous short-range dependence for $H < \frac{1}{2}$.

In particular, this behavior is induced only by the noise.

For $H > \frac{1}{2}$, if we replace hypothesis (ii)' with
 (ii)''

$$\lim_{s \rightarrow +\infty} s^{2-2H} \int_0^t e^{\frac{u}{\theta}} c(u, s) du = +\infty$$

we have again a time non-homogeneous long-range dependence,
 but this time

this behavior is induced by the covariance of the process I .

For $H > \frac{1}{2}$, if we replace hypothesis (ii)' with
 (ii)''

$$\lim_{s \rightarrow +\infty} s^{2-2H} \int_0^t e^{\frac{u}{\theta}} c(u, s) du = +\infty$$

we have again a time non-homogeneous long-range dependence,
 but this time

this behavior is induced by the covariance of the process I .

For $H < \frac{1}{2}$ we cannot conclude the same assertion.

The fractional Ornstein-Uhlenbeck process (fOU)

The fOU process with forcing term (ffOU)

The time-changed fractional Ornstein-Uhlenbeck process (TCfOU)

The limit distribution of the TCfOU process

Convergence results for the TCfOU process

The mean value function

The covariance function

Covariance as the function of time and Hurst index

The fact that V preserves its correlation for long times when $H > \frac{1}{2}$, may be useful in the field of neuronal modeling when one wants to include memory effects.

Now, we study the time-changed fractional OU (TCfOU) process and some convergence results concerning its one-dimensional distribution.

In particular, here, we show results concerning that,

- despite the time change, the process admits a Gaussian limit random variable.
- We prove that the process converges towards the time-changed OU (TCOU) as the Hurst index $H \rightarrow 1/2^+$, with locally uniform convergence of one-dimensional distributions.
- We also achieve convergence in the Skorohod J_1 -topology of the TCfOU process as $H \rightarrow 1/2^+$ in the space of càdlàg functions.

The fractional Ornstein-Uhlenbeck process (again)

In order to put in evidence the dependence on H , we adopt a slightly different notation and we consider again the fractional OU (fOU) process as the solution of the fractional Brownian motion (fBm)-driven equation

$$dU_H(t) = -\frac{1}{\theta} U_H(t) dt + \sigma dB_H(t),$$

where $\theta, \sigma \in \mathbb{R}^+$ and $B_H = \{B_H(t), t \geq 0\}$ is a fBm with Hurst parameter $H \in (0, 1)$.

Considering $H > 1/2$, we focus on the case of its long-range dependence.

The time-change

In what follows we will consider subordinators, i.e. increasing and starting from zero (hence positive) Lévy processes (see [Bertoin, 1996]). Each subordinator admits a Laplace exponent $\Phi(\lambda)$, i.e. a function $\Phi : [0, +\infty) \rightarrow \mathbb{R}$ such that

$$\mathbb{E}[e^{-\lambda\sigma(t)}] = e^{-t\Phi(\lambda)}, \quad t \geq 0, \lambda \geq 0.$$

In particular, such Laplace exponents Φ are Bernstein functions (see [Schilling, Song and Vondracek, 2012]) and therefore we can define the characteristic triplet of Φ , given by $(a_\Phi, b_\Phi, \nu_\Phi)$, where $a_\Phi, b_\Phi \geq 0$ are constants and ν_Φ is a Lévy measure on $(0, +\infty)$ such that

$$\int_0^{+\infty} (t \wedge 1) \nu_\Phi(dt) < +\infty.$$

Indeed, let us recall that any Bernstein function Φ can be represented in a unique way by means of the characteristic triple as

$$\Phi(\lambda) = a_\Phi + b_\Phi \lambda + \int_0^{+\infty} (1 - e^{-t\lambda}) \nu_\Phi(dt).$$

Vice versa, each Bernstein function Φ determines a unique (non-killed when $a_\Phi = 0$) subordinator $\sigma_\Phi(t)$ whose Laplace exponent is Φ .

Here we will consider Φ to be such that $a_\Phi = 0$ and if $b_\Phi = 0$ then we assume that $\nu_\Phi(0, +\infty) = +\infty$. Now let us define the inverse subordinator associated to Φ as

$$E_\Phi(t) := \inf\{y > 0 : \sigma_\Phi(y) > t\}.$$

It was established in [Meerschaert and Scheffler,, 2008] that our hypotheses on Φ are enough to guarantee that $E_\Phi(t)$ admits a probability density function $f_\Phi(s; t)$ for each $t > 0$.

A particular case is given by the choice $\Phi(\lambda) = \lambda^\alpha$ for $\alpha \in (0, 1)$.
Indeed, in this case, we have the α -stable subordinator $\sigma_\alpha(t)$.
According to [Meerschaert and Straka, 2013], $\sigma_\alpha(t)$ and $E_\alpha(t)$ are
absolutely continuous random variables for any $t > 0$.

Let us also fix some number $\theta > 0$ and focus on the fractional Ornstein-Uhlenbeck process as

$$U_H(t) = e^{-\frac{t}{\theta}} \int_0^t e^{\frac{s}{\theta}} dB^H(s), \quad t \geq 0, \quad (14)$$

which is a Gaussian process with one-dimensional density

$$p_H(t, x) = \frac{1}{\sqrt{2\pi V_{2,H}(t)}} e^{-\frac{x^2}{2V_{2,H}(t)}}, \quad x \in \mathbb{R}, \quad t \geq 0, \quad (15)$$

where

$$V_{2,H}(t) = e^{-2\frac{t}{\theta}} \int_0^t \int_0^t e^{\frac{v+u}{\theta}} |u - v|^{2H-2} du dv, \quad t \geq 0$$

is the **variance** of $U_H(t)$. Let us also denote by

$V_{n,H}(t) = \mathbb{E}[|U_H(t)|^n]$ the n th absolute moment of $U_H(t)$.

We construct the time-changed fractional Ornstein-Uhlenbeck process by considering a fractional Ornstein-Uhlenbeck process $U_H(t)$, together with an independent inverse subordinator $E_\Phi(t)$, and defining

$$U_{H,\Phi}(t) := U_H(E_\Phi(t)).$$

Let us state some properties of $V_{n,H,\Phi}(t) := \mathbb{E}[|U_{H,\Phi}(t)|^n]$.

Proposition

- (i) $V_{2n,H,\Phi}(t)$ is finite for any $t > 0$ and $n \in \mathbb{N}$.
- (ii) It holds that

$$V_{2n,H,\Phi}(t) = \int_0^{+\infty} V_{2n,H}(s) f_{\Phi}(s, t) ds.$$

- (iii) $V_{2n,H,\Phi}(t)$ is increasing in t for any $n \in \mathbb{N}$ and

$$\lim_{t \rightarrow +\infty} V_{2n,H,\Phi}(t) = V_{2n,H}(\infty) = \frac{(2\theta^{2H} H \Gamma(2H))^n \Gamma\left(\frac{2n+1}{2}\right)}{\sqrt{\pi}}.$$

Lemma

Function $V_{2,H}$ satisfies the relations

$$\lim_{t \rightarrow 0^+} \frac{V_{2,H}(t)}{t^{2H}} = 1 \text{ and } \lim_{t \rightarrow +\infty} V_{2,H}(t) = \theta^{2H} H \Gamma(2H).$$

Moreover, $V_{2,H} \in C^1(0, +\infty)$ and its derivative satisfies the relations

$$\lim_{t \rightarrow 0^+} \frac{V'_{2,H}(t)}{t^{2H-1}} = 2H \text{ and } \lim_{t \rightarrow +\infty} e^{\frac{t}{\theta}} t^{2-2H} V'_{2,H}(t) = 2H(2H-1)\theta.$$

We first focus on the one-dimensional convergence of $U_H(t)$ to the (classical) OU process $U(t)$, i.e.

$$U_H \rightarrow U \text{ (in distribution) as } H \rightarrow 1/2^+.$$

Recalling that $U_H(0) = 0$ and $\mathbb{E}[U_H(t)] = 0$, about the convergence of the variance and that of all the moments we have:

Proposition

For any $n \geq 1$ it holds that $\lim_{H \rightarrow \frac{1}{2}^+} V_{n,H}(t) = V_{n,\frac{1}{2}}(t)$ and the convergence is uniform in $[0, +\infty)$.

Now let us observe that, since $\mathbb{E}[U_H(t)] = 0$, $V_{2,H}(t) \rightarrow V_{2,\frac{1}{2}}(t)$, and $U_H(t)$ is a Gaussian random variable, we have that, for fixed $t > 0$,

$$U_H(t) \xrightarrow{d} U_{\frac{1}{2}}(t).$$

Thus we already have the weak one-dimensional convergence of the fractional Ornstein-Uhlenbeck process to the classical one as $H \rightarrow 1/2^+$. However, we can improve this kind of convergence, showing that the density $p_H(x, t)$ converges uniformly to $p_{1/2}(x, t)$.

Theorem

It holds that

$$\lim_{H \rightarrow \frac{1}{2}^+} p_H(t, x) = p_{\frac{1}{2}}(t, x)$$

for any $t \in (0, +\infty)$ and for any $x \in \mathbb{R}$.

The limit distribution of the TCfOU process

Now, let us explore the limit distribution of the TCfOU process as $t \rightarrow +\infty$.

It is well known that (for negative drift parameters) the fractional Ornstein-Uhlenbeck process is ergodic and admits Gaussian limit distribution with density function

$$p_H(\infty, x) = \frac{1}{\sqrt{2\pi V_H}} e^{-\frac{x^2}{2V_H}}$$

where $V_H := V_{2,H}(\infty)$.

By using this result we can exploit the limit distribution of the TCfOU process, i.e.

$$TCfOU \rightarrow fOU$$

for the time-limit distribution.

Proposition

Let $p_{H,\Phi}(t, x)$ be the probability density function of $U_{H,\Phi}(t)$. Then it holds that

$$\lim_{t \rightarrow +\infty} p_{H,\Phi}(t, x) = p_H(\infty, x). \quad (16)$$

Moreover, $U_{H,\Phi}(t) \xrightarrow{d} Z$ as $t \rightarrow +\infty$, where $Z \sim \mathcal{N}(0, V_H)$.

Let us observe that if we are able to establish equality (16), then the weak convergence result directly follows.

The proof relies on the following facts:



$$p_{H,\Phi}(t, x) = \int_0^{+\infty} p_H(s, x) f_\Phi(s; t) ds = \mathbb{E}[p_H(E_\Phi(t), x)], \quad (17)$$

where p_H is defined by equality (15),



$$p_H(E_\Phi(t), x) \leq \frac{1}{\sqrt{2\pi E_\Phi(1)}}. \quad (18)$$

valid for $t > 1$ being $E_\Phi(t)$ a.s. non-decreasing,

- $\mathbb{E} \left[\frac{1}{\sqrt{V_{2,H}(E_\Phi(1))}} \right] < +\infty$

- the use dominated convergence theorem to conclude that limit relation (16) holds, as $E_\Phi(t) \rightarrow +\infty$ as $t \rightarrow +\infty$ almost surely (being $a_\Phi = 0$ by assumption).

The fractional Ornstein-Uhlenbeck process (fOU)

The fOU process with forcing term (ffOU)

The time-changed fractional Ornstein-Uhlenbeck process (TCfOU)

The limit distribution of the TCfOU process

Convergence results for the TCfOU process

This result was expected. Indeed, the action of the time-change consists in change of the time scale that generally does not affect the limit distributions.

The fractional Ornstein-Uhlenbeck process (fOU)

The fOU process with forcing term (ffOU)

The time-changed fractional Ornstein-Uhlenbeck process (TCfOU)

The limit distribution of the TCfOU process

Convergence results for the TCfOU process




Figure: The density function $p_{H, \Phi}(t, x)$ for $\Phi(\lambda) = \sqrt{\lambda}$, $H = 3/4$, $\theta = 1$ and different values of t . The dashed line represents the limit distribution.

The fractional Ornstein-Uhlenbeck process (fOU)

The fOU process with forcing term (ffOU)

The time-changed fractional Ornstein-Uhlenbeck process (TCfOU)

The limit distribution of the TCfOU process

Convergence results for the TCfOU process

Weak one-dimensional convergence of the TCfOU process

A functional limit theorem for the TCfOU process

Weak one-dimensional convergence of the TCfOU process, $TCfOU \rightarrow TCOU$

Now we want to discuss the one-dimensional convergence of the TCfOU process to the time-changed Ornstein-Uhlenbeck process, i.e.

$$U_{H,\Phi} \rightarrow U_{\Phi}.$$

To do this, let us actually demonstrate some more strong convergence of the probability density function.

Theorem

Let $p_{H,\Phi}(t, x)$ be the probability density function of $U_{H,\Phi}(t)$ and $p_{\frac{1}{2},\Phi}(t, x)$ the probability density function of $U_{\Phi}(t)$. Then it holds

$$\lim_{H \rightarrow \frac{1}{2}^+} p_{H,\Phi}(t, x) = p_{\frac{1}{2},\Phi}(t, x) \quad (19)$$

for any $t > 0$ and $x \in \mathbb{R}$. Moreover, for any compact set $K \subset \mathbb{R} \setminus \{0\}$ it holds $p_{H,\Phi} \rightarrow p_{\frac{1}{2},\Phi}$ uniformly in $[0, +\infty) \times K$ as $H \rightarrow \frac{1}{2}^+$.

The fractional Ornstein-Uhlenbeck process (fOU)

The fOU process with forcing term (ffOU)

The time-changed fractional Ornstein-Uhlenbeck process (TCfOU)

The limit distribution of the TCfOU process

Convergence results for the TCfOU process

Weak one-dimensional convergence of the TCfOU process

A functional limit theorem for the TCfOU process




Figure: The density function $p_{H,\Phi}(t, x)$ for $\Phi(\lambda) = \sqrt{\lambda}$, $t = 1$, $\theta = 1$ and different values of H . The dashed line represents the case $H = 1/2$.

The fractional Ornstein-Uhlenbeck process (fOU)

The fOU process with forcing term (ffOU)

The time-changed fractional Ornstein-Uhlenbeck process (TCfOU)

The limit distribution of the TCfOU process

Convergence results for the TCfOU process

Weak one-dimensional convergence of the TCfOU process

A functional limit theorem for the TCfOU process

As for the fractional Ornstein-Uhlenbeck process, we can establish also the uniform convergence of the absolute moments.

Proposition

It holds that $V_{n,H,\Phi} \rightarrow V_{n,\frac{1}{2},\Phi}$ as $H \rightarrow \frac{1}{2}^+$ uniformly in $[0, +\infty)$.

A functional limit theorem for the TCfOU process

Let us consider the function space $C = C(0, +\infty)$ equipped with the uniform norm. The processes

$$U_{H,\Phi} = \{U_{H,\Phi}(t), t \geq 0\} \text{ (TCfOU),}$$

$$U_{\Phi} = \{U_{\Phi}(t), t \geq 0\}, \text{ (TCOU)}$$

$$U_H = \{U_H(t), t \geq 0\} \text{ (fOU),}$$

$$U = \{U(t), t \geq 0\} \text{ (OU),}$$

can be seen as C -valued random variables and then we can study the weak convergence in C of such random variables.

Let us denote by C^* the dual space of C (i.e. the space of continuous linear functionals on C).

For a sequence of C -valued random variables $(X_n)_{n \in \mathbb{N}}$ and a C -valued random variable $X \in C$, it holds

$$X_n \Rightarrow X$$

if and only if for any $\mathcal{F} \in C^*$ it holds $\mathbb{E}[\mathcal{F}(X_n)] \rightarrow \mathbb{E}[\mathcal{F}(X)]$ as $n \rightarrow +\infty$.

In particular, for any $\gamma \in (0, 1]$, let us denote by $\text{Lip}_\gamma([0, T])$ the space of Hölder-continuous functions on $[0, T]$ with exponent γ , which is a Banach space when equipped with the norm

$$\|f\|_{\text{Lip}_\gamma([0, T])} = \sup_{\substack{(t,s) \in [0, T]^2 \\ t \neq s}} \frac{|f(t) - f(s)|}{|t - s|^\gamma} + \|f\|_{L^\infty(0, T)}.$$

At first, let us focus on U_H and U .

Theorem

It holds $U_H \Rightarrow U$ in C as $H \rightarrow \frac{1}{2}^+$. Moreover, for any $T > 0$ and any linear functional $\mathcal{F} \in \text{Lip}_\gamma^*([0, T])$ for $\gamma < 1/2$ it holds $\mathbb{E}[\mathcal{F}(U_H)] \rightarrow \mathbb{E}[\mathcal{F}(U)]$.

The fractional Ornstein-Uhlenbeck process (fOU)

The fOU process with forcing term (ffOU)

The time-changed fractional Ornstein-Uhlenbeck process (TCfOU)

The limit distribution of the TCfOU process

Convergence results for the TCfOU process

Weak one-dimensional convergence of the TCfOU process

A functional limit theorem for the TCfOU process

Remark

The bound $\gamma < 1/2$ is sharp since U belongs to $\text{Lip}_\gamma([0, T])$ for $\gamma < 1/2$ but not for $\gamma = 1/2$.

Now we need to provide a similar result for

$$U_{H,\Phi} \text{ and } U_{\Phi}.$$

With this purpose, let us consider the set D of cadlag functions on $[0, +\infty)$ and Λ the set of strictly increasing functions $g : [0, +\infty) \rightarrow [0, +\infty)$. Let $\iota \in \Lambda$ be the identity function on $[0, +\infty)$.

The Skorohod J_1 metric on D is, for any couple of functions $f_1, f_2 \in D$,

$$d_{J_1}(f_1, f_2) = \inf_{g \in \Lambda} \max\{\|f_1 \circ g - f_2\|_{L^\infty(0, +\infty)}, \|g - \iota\|_{L^\infty(0, +\infty)}\}.$$

With this metric, the set (D, d_{J_1}) is a metric space (that we will denote only as D), thus we can consider the notion of weak convergence of D -valued random variables, which is actually weaker than the weak convergence in C .

With this in mind, let us show the following result that relies on a continuous mapping technique which is common in such kind of functional limit theorems (see, for instance, [Meerschaert and Scheffler, 2008, Scalas and Viles, 2014]).

Theorem

It holds $U_{H,\Phi} \Rightarrow U_\Phi$ in D as $H \rightarrow \frac{1}{2}^+$.

Proof.

First, note that C is a closed subspace of D , hence D^* is contained in C^* .

From $U_H \Rightarrow U$ in C we have the same convergence in D .

Consider the coupled processes (U_H, E_Φ) and (U, E_Φ) .

Since E_Φ is independent of both U_H and U , we get

$(U_H, E_\Phi) \Rightarrow (U, E_\Phi)$ in $D \times D$.

We remark that $U \in C$ almost surely and $E_\Phi \in D_\uparrow$ almost surely, where $D_\uparrow = \{f \in D : f \text{ is increasing}\}$.

Now let us denote by g the composition map on $D \times D$, i.e. $g(x, y) = x \circ y \in D$, and with $\text{Disc}(g)$ the set of discontinuity points of g .

Proof (Cont.)

It follows from [Whitt,2002] that

$$\text{Disc}(g) \subseteq (D \times D) \setminus ((C \times D_{\uparrow}) \cup (D \times C_{\uparrow\uparrow})),$$

where $D_{\uparrow\uparrow} = \{f \in D : f \text{ is strictly increasing}\}$ and $C_{\uparrow\uparrow} = C \cap D_{\uparrow\uparrow}$. Taking into account the inclusion $(U, E_{\Phi}) \in C \times D_{\uparrow}$ almost surely, we get that

$$\mathbb{P}((U, E_{\Phi}) \in \text{Disc}(g)) = 0.$$

Thus we can use the continuous mapping theorem to obtain the desired convergence. □

-  Kaarakka T, Salminen P (2011) On fractional Ornstein-Uhlenbeck processes. *Communications on Stochastic Analysis* 5(1):8, DOI 10.31390/cosa.5.1.08
-  Cheridito P, Kawaguchi H, Maejima M (2003) Fractional Ornstein-Uhlenbeck processes. *Electronic Journal of Probability* 8, doi:10.1214/EJP.v8-125.
-  Biagini F., Hu Y., Øksendal B., Zhang T. (2008) Stochastic calculus for fractional Brownian motion and applications. Springer Science & Business Media
-  Borovkov K, Mishura Y, Novikov A, Zhitlukhin M (2017) Bounds for expected maxima of Gaussian processes and their discrete approximations. *Stochastics* 89(1):21–37, doi:10.1080/17442508.2015.1126282
-  Richard A., Orio P. & Tanré E., An integrate-and-fire model to generate spike trains with long-range dependence, *J. Comput. Neurosci.* (2018) 44: 297.



Ascione G., Mishura Y., P. E., Fractional Ornstein-Uhlenbeck process with stochastic forcing and its applications, Methodology and Computing in Applied Probability, 2021, 23(1), pp. 53–84.



Ascione G., Mishura Y., P. E., Time-changed fractional Ornstein-Uhlenbeck process, Fractional Calculus and Applied Analysis 23, 2020, no. 2, 450–483.



Ascione G., Mishura Y., P. E., Convergence results for the time-changed fractional ornstein–uhlenbeck processes, Theory of Probability and Mathematical Statistics, 2021, 104, pp. 23–47.



Ascione G., Mishura Y., P. E., The Fokker-Planck equation for the time-changed fractional Ornstein-Uhlenbeck process, Proceedings of the Royal Society of Edinburgh Section A: Mathematics, 2021.



Mishura Y (2008) Stochastic calculus for fractional Brownian motion and related processes, vol 1929. Springer Science and Business Media

The fractional Ornstein-Uhlenbeck process (fOU)

The fOU process with forcing term (ffOU)

The time-changed fractional Ornstein-Uhlenbeck process (TCfOU)

The limit distribution of the TCfOU process

Convergence results for the TCfOU process

Weak one-dimensional convergence of the TCfOU process

A functional limit theorem for the TCfOU process

THANK YOU FOR YOUR KIND ATTENTION