

## Stochastic wave equations with super-linear coefficients

Marta Sanz-Solé, Universitat de Barcelona and BGSMath

We consider a stochastic wave equation on  $\mathbb{R}^d$ ,  $d \in \{1, 2, 3\}$ , driven by a Gaussian noise in  $(t, x)$ , white in time. We assume that the free terms  $b$  and  $\sigma$  are such that, for  $|x| \rightarrow \infty$ ,

$$|\sigma(x)| \leq \sigma_1 + \sigma_2|x|(\ln_+(|x|))^a, \quad |b(x)| \leq \theta_1 + \theta_2|x|(\ln_+(|x|))^\delta, \quad (1)$$

where  $\theta_2, \sigma_2 > 0$ ,  $\delta, a > 0$ , with  $b$  *dominating* over  $\sigma$ . For any fixed time horizon  $T > 0$  and with a suitable constraints on the parameters  $a$ ,  $\delta$ ,  $\sigma_2$  and  $\theta_2$ , we prove existence of a random field solution to the equation and that this solution is unique, and bounded in time and in space a.s.

The research is motivated by the article [R. Dalang, D. Khoshnevisan, T. Zhang, *AoP*, 2019] on a 1-d reaction-diffusion equation with coefficients satisfying conditions similar to (1). We see that the  $L^\infty$ - method used by these authors can be successfully implemented in the case of wave equations. This is joint work with A. Millet (U. Paris 1, Panthéon-Sorbonne).