

A characterization of balanced rational normal surface scrolls in terms of their osculating spaces II

EDOARDO BALLICO, RAGNI PIENE, AND HSIN-SHENG TAI¹

In this note we prove a result conjectured by the second and third author [3, Conjecture (i), p. 216]; the proof was found independently by the first author and by the second and third authors. The results needed in the proof are contained in [3]; it only remains to apply a theorem of Van de Ven [6, Theorem III, p. 406] to the adjunction divisor. We would like to take this opportunity to point out adjunction theory as a very powerful tool in projective geometry!

Let $X \subset \mathbf{P}^N$ be a smooth, complex, projective surface. Recall [3, p. 216] that the m th osculating space to X at a point x is the linear subspace $\text{Osc}_X^m(x)$ determined by the partial derivatives of order $\leq m$ of the coordinate functions, with respect to a system of local parameters for X at x and evaluated at x . More precisely, let $\mathcal{P}_X^m(1)$ denote the sheaf of principal parts of order m of $\mathcal{O}_X(1) := \mathcal{O}_{\mathbf{P}^N}(1)|_X$. Then $\mathcal{P}_X^m(1)$ is locally free with rank $\binom{m+2}{2}$, and there are homomorphisms

$$a^m: \mathcal{O}_X^{N+1} \rightarrow \mathcal{P}_X^m(1)$$

such that $\text{Im}(a^m(x))$ defines the m -th order osculating space to X at x , i.e., such that

$$\text{Osc}_X^m(x) := \mathbf{P}(\text{Im}(a^m(x))) \subset \mathbf{P}^N.$$

For a general surface, one expects the dimension of $\text{Osc}_X^m(x)$ to be $\binom{m+2}{2} - 1$ for almost all points x of X and all m such that $\binom{m+2}{2} - 1 \leq N$. Points where this dimension is smaller than expected are “flat” points of the surface — often called points of hyperosculation. If the surface X contains a line through the point x , then the dimension of $\text{Osc}_X^m(x)$ is at most $2m$. Hence a ruled surface has the property that all its m th order osculating spaces have dimension at most $2m$. There are, however, non-ruled surfaces with this property (Togliatti [5], Dye [1]).

Suppose the surface X is a *balanced rational normal scroll of degree $2n$* , i.e., X is isomorphic to $\mathbf{P}^1 \times \mathbf{P}^1$ embedded in $\mathbf{P}^{2n+1}$ via $\text{pr}_1^* \mathcal{O}_{\mathbf{P}^1}(1) \otimes \text{pr}_2^* \mathcal{O}_{\mathbf{P}^1}(n)$. It is shown in [2, p. 1060] that in this case all m th order osculating spaces to X have dimension $2m$, for $m \leq n$. The theorem we shall prove shows that this property characterizes the balanced rational normal scrolls.

¹Supported in part by the Norwegian Research Council for Science and the Humanities and by the National Nature Science Foundation of China (Tianyuan 19136003).

1980 *Mathematics subject classifications*. Primary 14J26; Secondary 14F10, 14M12, 14N05.

THEOREM. *Let $X \subset \mathbf{P}^{2n+1}$ be a smooth, projective surface, not contained in a hyperplane, such that $\dim \text{Osc}_X^m(x) = 2m$ for all $x \in X$ and all $m \leq n$. Then X is a balanced rational normal scroll of degree $2n$.*

PROOF. In [3, Theorem, p. 221] the second and third author proved that any surface satisfying the hypotheses of the theorem is birationally ruled, but not isomorphic to $\mathbf{P}^2$. Moreover, the theorem was shown to hold if X is geometrically ruled (i.e., has no exceptional curves of the first kind) or if X is rational. Finally, it was shown that the theorem holds provided $n \leq 4$.

We shall prove the theorem by showing that the assumptions $n \geq 4$, X is not rational, X is birationally ruled and contains at least one exceptional curve, lead to a contradiction.

Let H denote the class of a hyperplane section of X and let K denote the class of a canonical divisor on X . Since X spans $\mathbf{P}^{2n+1}$, we have

$$\dim H^0(X, \mathcal{O}_X(1)) \geq 2n + 2 \geq 12$$

and $\deg X = H^2 \geq 2n \geq 10$. By a theorem of Van de Ven [6, Theorem III, p. 406] it follows that $K + H$ is very ample unless there exists an exceptional curve E on X with $E.H = 1$. Let us show that such an E cannot occur. Recall from [3, p. 220] that X possesses a line bundle $\mathcal{Q}$, given by

$$\mathcal{Q} := \text{Coker}(a^2: \mathcal{O}_X^{N+1} \rightarrow \mathcal{P}_X^2(1)),$$

with first Chern class

$$c_1(\mathcal{Q}) = \frac{1}{n-1}(2nK + (n+3)H).$$

Since $E.K = -1$, $E.H = 1$ implies

$$c_1(\mathcal{Q}).E = -\frac{n-3}{n-1}.$$

But this is not an integer since we have assumed $n \geq 5$. So $K + H$ must be very ample.

We can therefore argue as in [3, p. 221], using $K + H$ instead of H . The theorem of Sommese [4, Theorem (1.5), p. 377] and Van de Ven [6, Theorem II, p. 403] allows us to conclude that $2K + H$ is generated by its global sections, in particular that $(2K + H)^2 \geq 0$ holds.

Let $\tau = \frac{1}{3}(K^2 - 2c_2(X))$ denote the Hirzebruch index of X . Since X is birationally ruled and not isomorphic to $\mathbf{P}^2$, we have $K^2 = 8(1 - q) - t$ and $c_2(X) = 4(1 - q) + t$, where t denotes the number of exceptional curves on X with respect to a relatively minimal model, and q denotes the genus of the base curve of this minimal model. Therefore $\tau = -t$ holds, hence $\tau \leq 0$.

Recall [3, (2'), p. 220; p. 221] that the numerical characters of X satisfy the formulas

$$n(2n+1)K^2 + 2n(n+5)K \cdot H + 2(n^2 + 2n + 3)H^2 = 0$$

and

$$3n(2n+1)\tau = -2(n+3)(nK \cdot H + (n+1)H^2).$$

From these two formulas we deduce the following equality

$$(2K+H)^2 = \frac{54}{n+3}\tau - 3\frac{n-4}{n}H^2.$$

Since $\tau \leq 0$, $n \geq 5$, and $H^2 > 0$, we get $(2K+H)^2 < 0$, which is the desired contradiction.

REFERENCES

- [1] R. H. Dye, *The extraordinary higher tangent spaces of certain quadric intersections*, Proc. Edinburgh Math. Soc. (to appear) x (199x), xx-xx.
- [2] R. Piene and G. Sacchiero, *Duality for rational normal scrolls*, Comm. Alg. 12(9) (1984), 1041-1066.
- [3] R. Piene and H. Tai, *A characterization of balanced rational normal scrolls in terms of their osculating spaces*, in "Enumerative Geometry, Proc. Sitges 1987," S. Xambó-Descamps (ed.), Lecture Notes in Math. 1436, Springer-Verlag, 1990, pp. 215-224.
- [4] A. J. Sommese, *Hyperplane sections of Projective surfaces I — the adjunction mapping*, Duke Math. J. 46 (1979), 377-401.
- [5] E. Togliatti, *Alcuni esempi di superficie algebriche degli iperspazi che rappresentano un'equazione di Laplace*, Comm. Math. Helv. 1 (1929), 255-272..
- [6] A. Van de Ven, *On the 2-connectedness of very ample divisors on a surface*, Duke Math. J. 46 (1979), 403-407.

Dipartimento di Matematica, Università degli studi di Trento, 38050 Povo, Trento, Italia
 Matematisk Institutt, Universitetet i Oslo, Box 1053 Blindern, 0316 Oslo 3, Norway
 Institute of Mathematics, Academia Sinica, Beijing 100080, China

